Title: Potato Disease Management – 2010 Proposal

Investigators: Robert D. Davidson and Andrew J. Houser

Department of Horticulture and Landscape Architecture

San Luis Valley Research Center

Funding requested from: CPAC

# Nature and Objectives:

This is the annual general request for funds to support approximately 50% of the cost of a range of research projects related to control and management of potato diseases specific to the San Luis Valley. For the most part, these are long term projects designed to address evolving chronic disease problems which limit potato production. Funds being requested are for fixed expenses: primarily labor (full time and hourly) and land rent. Each year the scope of work accomplished within this project expands, however, the base level funding request has remained relatively flat.

The potato industry in the San Luis Valley is constantly faced with serious disease management challenges. These challenges have been made more difficult recently because of the water situation and the potential for new strains of some of the more serious disease issues. Seed borne diseases such as powdery scab, pink rot, and silver scurf require increasingly sophisticated management schemes. New problems such as Zebra chip continue to be potential threats to the Colorado potato industry. While growers seem to have a handle on control of the foliar phase of early blight, the tuber phase continues to be damaging and difficult to control. Potato late blight was found in the Valley in 2007, but was eradicated in 2008 by the SLV producers through the use of well timed fungicide applications, seed lot screening and numerous grower education programs. Post harvest test plots for certified seed as well as field inspections indicate that potato leaf roll levels are down significantly and represent a fairly low incidence in the overall crop. Potato virus Y, however, is now epidemic in many cultivars, particularly Russet Norkotah. In the short term, certified seed stocks of several cultivars will be quite limited. The proposed Certified Seed Law for Colorado will go a long ways toward reducing inoculum, but will not be highly effective in the short term. Instead, the potato industry will need to flush out the high mosaic stocks and consider the use of more resistant cultivars to replace the acreage of these more problematic ones. It is of note that the new cultivar releases from the Colorado cultivar development program are being screened for their reaction and symptomology to PVY and have performed quite well even during this serious epidemic.

Efficacy trials are a permanent component of this overall research effort. These trials generate the basic information required by product manufacturers, the EPA and State Department of Agriculture for labeling and registration. Ongoing evaluation of products is essential to maintaining current labels as well as justifying Section 18, LSN24C, and Section 3 registrations for new products or established products with label variations. Availability of new crop care products to potato growers in Colorado is contingent upon scientifically valid data developed within the University on a regional basis. Resistance management among fungicides, both foliar and soil applied is becoming even more critical as many products have similar modes of action. Therefore, development of comprehensive season-long fungicide programs for potato which rotate various chemistries with different modes of action are requiring more attention than in the past. There has been a steady movement of more of these types of trials to a controlled greenhouse environment due to the inability to accurately predict how various field trials will develop.

Finally, evaluation of advanced clones from the CSU Cultivar Development program for reaction to several critical diseases will continue as a permanent component of this research effort. The increasing threat of major diseases, especially from out-of-Valley sources, and the use of Plant Variety Protection for new cultivars make this work central to protecting grower and University interests. In addition, these evaluations provide valuable information to the producer of new cultivars so that the threat from many diseases is mitigated. A real emphasis will be placed this year and in the future in developing complete informational management sheets for each new cultivar that are grower friendly and useable. Disease management summaries will be formulated which outline the best management guidelines currently available for controlling specific diseases with emphasis on the practical grower tools available.

#### Methods, Procedures and Facilities:

Objective 1: Efficacy trials: Trials will occur at the SLV Research Center and the Selter's corner. Early blight, pink rot, silver scurf, and seed piece treatment/decay trials will be conducted on station under solid set irrigation. Early blight trials will continue to focus on the proper order for applying fungicides to maximize efficacy against early blight and late blight with consideration for appropriate rotations of the various chemicals and modes of action. Additional trials will be utilized to test new chemistries under SLV environmental conditions including the use of compost type products.

Objective 2: Advanced selection disease evaluations: Evaluation of advanced selections from the CSU Cultivar Development program for reactions to PLRV, PVY, and TRV as well as reactions to ring rot will be conducted off-station on the Selter's corner. Evaluation for powdery scab, PVY, early blight tuber rot, dry rot, and soft rot will be conducted in the SLVRC Processing building or the isolation greenhouses. This portion of the project will be in cooperation with Dr. David Holm.

Objective 3: Powdery scab management: Screening compounds for activity against Spongospora subterranea will be conducted at the station in the isolation greenhouses. Focus will center on development of an effective management program which will use soil screening for spore load (using PCR performed at the SLRVC), cultivar rotation, irrigation management, and use of Omega (fluazinam). Development of the appropriate delivery system and nozzle package for placement of Omega fungicide to control powdery scab will be finalized. Additionally, screening environmentally friendly chemistries such as compost teas (as part of a larger EPA grant) will be conducted both on station and at off-station sites, if available. A comprehensive greenhouse study is underway to examine the role of cultivar rotation in increasing or reducing the soil spore load. Root galls have been examined to better understand their role in the increase of soil borne inoculum in cultivars which show tuber resistance to lesion development. This portion of the project will be in cooperation with Dr. Sastry Jayanty.

Objective 4: Biological agent trials: Evaluation of biological agents, primarily compost and compost teas, and products to increase plant defense systems and corresponding yield and quality will be evaluated on station. This portion of the project will be in cooperation with Dr. Samuel Essah and Merl Dillon (returning as an hourly hire to work with several research projects).

Objective 5: Calcium placement trials: Calcium formulations will be utilized during different times of the season to examine their control and management of the physiological disorder known as buck skinning which occurs often on many of the more popular specialty cultivars and red potatoes. The SLVRC plus one grower cooperator's field will be used in this effort. This portion of the project will be in cooperation with Dr. Samuel Essah.

Objective 6: BMP's for specific diseases: Best management practices based on extensive literature reviews are being formulated for potential management of the following diseases: pink rot, silver scurf, PVY, and corky ring spot. Zebra chip information will be discussed in a series of newsletter type articles. Current field inspections have not identified Zebra chip to be present in Colorado certified seed, but the pathogen has been linked to Colorado produced potatoes in the past. Cultivar management sheets as well as these BMP's will be posted to the SLVRC web site in addition to being available for local producers.

### Resource needs at the SLVRC:

All resources necessary to conduct these projects are currently available or will be purchased as needed.

# Relationship of Proposed Research to Overall Problem:

The proposed research is driven by the mix of disease problems that limit potato production and quality of the crop in the San Luis Valley. The specific components of this research effort have been identified and ranked by SLV potato growers through annual surveys and direct contacts.

# **Potential for Leveraging Outside Funding:**

Historically, funds granted by the CPAC to the Potato Disease Management research project have been used to leverage outside funds from agricultural chemical companies, Colorado Certified Potato Growers' Association, CSU, AES, and USDA-CSREES. In general terms, for each dollar of base level funding from the CPAC, there have been two to three dollars leveraged from outside, non-Colorado potato industry, sources. It is, however, only by virtue of consistent base level funding that assures the presence of the project Research Associate, Andrew Houser, that these outside sources of funding can be secured.

### Time line for Proposed Research:

These projects are in various stages of completion. For efficacy trials, data are generated each year and used in registration and labeling of new products and local use recommendations for existing and new products. Data from cultivar evaluation studies is derived year-to-year and used by the CSU Cultivar Development program and states in the Western Regional Cultivar Development Program in the long term assessment of new releases, development of cultural management sheets for each cultivar released, and in preparing applications for Federal Plant Variety Protection. Research on powdery scab was formally started in 2001 and was primarily focused on the field. This next phase of the project will be focused more on the greenhouse and understanding how the fungus populations change over time based on cultivars grown. This is expected to continue for the coming two years. Several approaches to control and management of this disease are being pursued. Studies on the control and biology of the pink rot fungus will continue with clonal evaluation for symptoms being a priority. The study of biologicals and compost teas will last one more year. Corky ring spot symptom expression in new clones started in field trials in 2009 (in cooperation with Agro Engineering and Dr. Russ Ingham) will be moved to the isolation greenhouses at the SLVRC.

# Progress in 2009:

- Confirmed the cost effectiveness and the ability to control foliar early blight disease when applying three fungicides at the proper time. Continued to verify that the 650 DD model used for early blight is accurate and should be an excellent starting point for fungicide applications. Demonstrated that fungicide programs involving two to three applications, with one of the applications being a strobilurin class fungicide (especially when utilized as a first application treatment), reduced disease to acceptable levels. Proved that use of fungicides over years does not only significantly reduce the threat of disease, but also increases the yield of potatoes. Demonstrated to growers that as use of fungicides has increased, the separation between the untreated controls and the treated plots has reduced, indicating that there is a better opportunity for organic producers to maintain their crops with no fungicide treatments and that commercial production is seeing a benefit from fungicide application.
- Worked closely with the local consultants/field men, the Federal/State Inspection Service and growers to keep late blight out of the SLV potato crop. This included numerous meetings, seed lot screening, and having available the appropriate information (updated late blight management plan with current chemistries available for use) for growers. Two imported seed lots were confirmed with late blight in 2009 and were removed from planting in the SLV. Recorded late blight severity units at three valley locations (Blanca, Hooper, and Sargent), calculated and reported on the SLVRC web page and on phone message (at SLVRC) the early blight and nematode degree days throughout 2009 season to assist in timely, effective application of fungicides and/or insecticides.
- Confirmed the effectiveness of Omega in controlling powdery scab when applied in-furrow at planting. Confirmed that use of a split nozzle during application of in-furrow Omega (seed piece and surrounding soil) was more effective than use of only spraying over the seed piece. Earlier results from this project helped obtain a LSN24C label for commercial use of Omega on potatoes for the 2007 season. Overall grower success when using Omega as part of a complete management program has been excellent. Worked with compost and compost teas by themselves and in combination with Omega. Found that compost and compost teas by themselves did little to control powdery scab, but when using compost in combination with Omega, there was an increase in disease control. This will be evaluated further in 2010.
- Continued to work closely with seed growers on roguing principles for the cultivar Russet Norkotah and confirmed how mosaic symptoms appear in the field during the season, how effective is roguing, potential for

mechanical spread during the early season, and yield loss expected due to current season spread as compared with previous season's infection. Began documenting how a seed grower could best keep PVY to a minimal level through the use of several management options.

# Objectives and Expected Accomplishments for 2010:

- Supply a comprehensive data package on the disease reactions of all new potato clones released from CSU. Reduce potential for the release of problematic cultivars. Conduct earlier screening of new germplasm for resistance to powdery scab and potato virus Y in the greenhouse. Restructure and prioritize critical information necessary for growth of new and existing cultivars through the cultivar management sheets. Make the management sheets more users friendly and applicable to SLV grower's conditions.
- Continue work toward the development of a long term management strategies for pink rot and silver scurf in the SLV using existing research information and supplementing with focused research under SLV conditions. Continue to finalize how growers should raise PVY susceptible cultivars to produce stocks with the lowest levels of PVY increase possible. Best management practices for all phases of production will be formulated with the last several years of PVY research and available to growers through mailings and on the web.
- Finalize the development of a multi-component management regime for powdery scab for use where a determined degree of control is economically justified. Incorporate the research with the fungicide Omega; infurrow placement in the hill, use of appropriate nozzle packages, levels of control based upon soil potential, etc. Included in this effort is the development of long term management strategies to reduce potential build up of powdery scab in SLV soils by determining the best potato cultivars to use in rotation under various field scenarios. Continue use of the PCR test developed by Christ et al. for soil assays representing the potential spore population in the soils for local producers. Continue studies in a greenhouse setting to verify efficacy of certain chemistries as well as cultivar rotation in reducing powdery scab root galling and tuber symptoms.
- Generate information for potato growers, through efficacy trials, that can be used to reduce expenditures on fungicides based on specific data for crop care products addressing rates, combinations, sequences, rotations, season long programs, application timing, and less costly generic alternatives. Additionally, provide timely information about late blight potential in the Valley by continued cooperation in the effort to inspect all out-of-state seed sources, pursue more aggressive fungicide spray programs, and provide consistent communication with the overall industry during the season.
- Continue to evaluate biological chemistries, compost, compost teas, and other products to manage disease problems and/or increase plant defense systems and corresponding yield and quality.

Funding History: 2000, \$21,000; '01, \$18,000; '02, \$27,000; '03, \$25,000; '04, \$29,000; '05, \$20,000;

'06, \$25,000; '07, \$25,000; '08, \$30,000; '09, 33,000

Budget for 2010: \$33,000

Labor: Full time (50% Research Associate) + part time plot labor
Land rent (\$1,200) + supplies (\$3,800)

\$28,000
\$5,000